ELSEVIER

Neuroscience Letters 289 (2000) 235–238

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 289

Açikgöz, O., see Gönenç, S., 107
Abdullah, L.I., see Crawford, F.C., 61
Abe, H., see Watanabe, H., 53
Addicks, K., see Hess, A., 72
Aguilar, R., see González, D., 111
Aihara, Y., see Hasegawa, H., 161
Akhisaroglu, M., see Gönenç, S., 107
Aldudo, J., see Bullido, M.J., 213
Allikmets, L., see Matto, V., 131
Alonso, R., see González, D., 111
Annepu, J., Ravindranath, V., 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced complex I inhibition is reversed by disulfide reductant, dithiothreitol in mouse brain, 209
Aparecida Del Bel, E., see de Oliveira, R.M.W., 123
Artiga, M.J., see Bullido, M.J., 213

Bagli, M., see Majores, M., 224
Bai, D., see Sattelle, D.B., 197
Barbanel, G., see Laplanche, L., 49
Bayer, T.A., see Block, W., 147
Blasevich, M., see Petroni, A., 139
Bloch, W., see Hess, A., 72
Block, W., Bayer, T.A., Tepest, R., Träber, F., Rietschel, M., Müller, D.J., Schulze, T.G., Honer, W.G., Maier, W., Schild, H.H., Falkai, P., Decreased frontal lobe ratio of *N*-acetyl aspartate to choline in familial schizophrenia: a proton magnetic resonance spectroscopy study, 147
Boireau, A., Maréchal, P.-A., Meunier, M., Dubédat, P., Moussaoui, S., The anti-oxidant ebselen antagonizes the release of the apoptogenic factor cytochrome *c* induced by Fe^{2+}/citrate in rat liver mitochondria, 95
Bouët, V., Gahéry, Y., Muscular exercise improves knee position sense in humans, 143
Brownstein, M.J., see Charles, V., 29
Bullido, M.J., Guallar-Castillón, P., Artiga, M.J., Ramos, M.C., Sastre, I., Aldudo, J., Frank, A., Coria, F., Rodríguez-Artalejo, F., Valdivieso, F., Alzheimer's risk associated with human apolipoprotein E, alpha-2 macroglobulin and lipoprotein receptor related protein polymorphisms: absence of genetic interactions, and modulation by gender, 213
Burghoff, M., see Mackert, B.-M., 33

Carp, R.I., see Choi, Y.-G., 173
Casper, D., Yaparpalvi, U., Rempel, N., Werner, P., Ibuprofen protects dopaminergic neurons against glutamate toxicity in vitro, 201
Cavarretta, I., see Petroni, A., 139
Cecchi, C., see Gestri, D., 157
Charles, V., Mezey, E., Reddy, P.H., Dehejia, A., Young, T.A., Polymeropoulos, M.H., Brownstein, M.J., Tagle, D.A., Alpha-synuclein immunoreactivity of huntingtin polyglutamine aggregates in striatum and cortex of Huntington's disease patients and transgenic mouse models, 29
Chen, H.H., see Sattelle, D.B., 197
Cheong, J., see Moon, C., 41
Chey, J., Kim, J.-W., Cho, H.-Y., Effects of apolipoprotein E phenotypes on the neuropsychological functions of community-dwelling elderly individuals without dementia, 230
Cho, H.-Y., see Chey, J., 230
Choi, E.-K., see Choi, Y.-G., 173
Choi, K.-Y., see Won, M.H., 119
Choi, Y.-G., Kim, J.-I., Lee, H.-P., Jin, J.-K., Choi, E.-K., Carp, R.I., Kim, Y.-S., Induction of heme oxygenase-1 in the brains of scrapie-infected mice, 173
Clow, A., see Gallagher, I.M., 221
Copray, S., Kernell, D., Neurotrophins and TRK-receptors in adult rat spinal motoneurons: differences related to cell size but not to 'slow/fast' specialization, 217
Coria, F., see Bullido, M.J., 213
Crawford, F.C., Freeman, M.J., Schinka, J., Abdullah, L.I., Richards, D., Sevush, S., Duara, R., Mullan, M.J., The genetic association between Cathepsin D and Alzheimer's disease, 61
Cropper, E.C., see Klein, A.N., 37
Cunha, R.A., Ribeiro, J.A., Adenosine A_{2A} receptor facilitation of synaptic transmission in the CA1 area of the rat hippocampus requires protein kinase C but not protein kinase A activation, 127
Curio, G., see Mackert, B.-M., 33

Dabbert, D., Rösner, S., Krämer, M., Schöll, U., Tumani, H., Mäder, M., Weber, F., Glatiramer acetate (Copolymer-1)-specific, human T cell lines: cytokine profile and suppression of T cell lines reactive against myelin basic protein, 205
De Keyser, J., see Zeinstra, E., 75
de Oliveira, R.M.W., Aparecida Del Bel, E., Mamede-Rosa, M.L.N., Padovan, C.M., Deakin, J.F.W., Guimarães, F.S., Expression of neuronal nitric oxide synthase mRNA in stress-related brain areas after restraint in rats, 123
de Tommaso, M., Guido, M., Libro, G., Sciruicchio, V., Puca, F., Zolmitriptan reverses blink reflex changes induced during the migraine attack in humans, 57
Deakin, J.F.W., see de Oliveira, R.M.W., 123
Dehejia, A., see Charles, V., 29
Duara, R., see Crawford, F.C., 61
Dubédat, P., see Boireau, A., 95

Eidizadeh, S., see Pennypacker, K., 1
Eiselt, M., see Zwiener, U., 103

Faden, A.I., see Knoblach, S.M., 5
Falkai, P., see Block, W., 147
Fekete, C., Légrádi, G., Mihály, E., Tatro, J.B., Rand, W.M., Lechan, R.M.,

PII: S0304-3940(00)01331-8

α-Melanocyte stimulating hormone prevents fasting-induced suppression of corticotropin-releasing hormone gene expression in the rat hypothalamic paraventricular nucleus, 152
Flink, R., see Samuelsson, C., 185
Frank, A., see Bullido, M.J., 213
Freeman, M.J., see Crawford, F.C., 61
Fujii, T., Harada, H., Koyama, T., Nakajima, Y., Kawashima, K., Effects of physostigmine and calcium on acetylcholine efflux from the hippocampus of freely moving rats as determined by in vivo microdialysis and a radioimmunoassay, 181
Fujimoto, M., see Wang, X., 115

Gahéry, Y., see Bouët, V., 143
Gallagher, I.M., Jenner, P., Glover, V., Clow, A., CuZn-superoxide dismutase transgenic mice: no effect on longevity, locomotor activity and ^{3}H-mazindol and ^{3}H-spiperone binding over 19 months, 221
Galli, C., see Petroni, A., 139
Garrido-Gracia, J.C., see González, D., 111
Gestri, D., Cecchi, C., Tedde, A., Latorraca, S., Orlacchio, A., Grassi, E., Massaro, A.M., Liguri, G., St. George-Hyslop, P.H., Sorbi, S., Lack of SOD1 gene mutations and activity alterations in two Italian families with amyotrophic lateral sclerosis, 157
Ghafouri, M., Lestienne, F.G., Altered representation of peripersonal space in the elderly human subject: a sensorimotor approach, 193
Gießler, F., see Zwiener, U., 103
Glatzle, J., see Kreis, M.E., 45
Glover, V., see Gallagher, I.M., 221
Gönenç, S., Açikgöz, O., Kayatekin, B.M., Uysal, N., Akhisaroglu, M., Effects of footshock stress on superoxide dismutase and glutathione peroxidase enzyme activities and thiobarbituric acid reactive substances levels in the rat prefrontal cortex and striatum, 107
González, D., Bellido, C., Aguilar, R., Garrido-Gracia, J.C., Hernández, G., Alonso, R., Sánchez-Criado, J.E., Luteinizing hormone secretion elicited in a ligand-independent activation of progesterone receptor manner at pituitary level in the rat: differential effect of two selective estrogen receptor modulators, 111
Grassi, E., see Gestri, D., 157
Grundy, D., see Kreis, M.E., 45
Guallar-Castillón, P., see Bullido, M.J., 213
Guido, M., see de Tommaso, M., 57
Guimarães, F.S., see de Oliveira, R.M.W., 123

Harada, H., see Fujii, T., 181
Harrison, J.B., see Sattelle, D.B., 197
Hasebe, Y., see Nagai, M., 227
Hasegawa, H., Yazawa, T., Yasumatsu, M., Otokawa, M., Aihara, Y., Alteration in dopamine metabolism in the thermoregulatory center of exercising rats, 161
Hernández, G., see González, D., 111
Hess, A., Michel, O., Kopec, T., Wittekindt, C., Bloch, W., Stennert, E., Addicks, K., Detection of extracellular signal-regulated kinase1/2 in the inner ear of guinea pigs, 72
Heun, R., see Majores, M., 224
Hiss, L.-H., see Mackert, B.-M., 33
Honer, W.G., see Block, W., 147
Horikomi, K., see Takahashi, S., 21

Ichinose, H., see Mogi, M., 165
Ishida, Y., see Shirokawa, T., 69
Isobe, K.-i., see Shirokawa, T., 69
Ito, E., see Takagi, H., 169

Jee, Y., see Moon, C., 41
Jenner, P., see Gallagher, I.M., 221
Jeong, Y.G., see Won, M.H., 119
Jessen, F., see Majores, M., 224
Jin, J.-K., see Choi, Y.-G., 173
Jo, S.M., see Won, M.H., 119

Kaba, H., see Osako, Y., 25
Kaehler, S.T., see Singewald, N., 17
Kakigi, R., see Wang, Y., 135
Kakuda, T., Yanase, H., Utsunomiya, K., Nozawa, A., Unno, T., Kataoka, K., Protective effect of γ-glutamylethylamide (theanine) on ischemic delayed neuronal death in gerbils, 189
Kamenka, J.-M., see Laplanche, L., 49
Kang, T.-C., see Won, M.H., 119
Kapron, C.M., see Sylvester, L., 13
Kassed, C.A., see Pennypacker, K., 1
Kataoka, K., see Kakuda, T., 189
Kawashima, K., see Fujii, T., 181
Kayatekin, B.M., see Gönenç, S., 107
Kernell, D., see Copray, S., 217
Kim, H., see Moon, C., 41
Kim, J.-I., see Choi, Y.-G., 173
Kim, J.-W., see Chey, J., 230
Kim, S., see Moon, C., 41
Kim, Y.-S., see Choi, Y.-G., 173
Klein, A.N., Weiss, K.R., Cropper, E.C., Glutamate is the fast excitatory neurotransmitter of Small Cardioactive Peptide – containing *Aplysia* radula mechanoafferent neuron B21, 37
Knoblach, S.M., Faden, A.I., Cortical interleukin-1β elevation after traumatic brain injury in the rat: no effect of two selective antagonists on motor recovery, 5
Kopec, T., see Hess, A., 72
Kouvelas, D., see Singewald, N., 17
Koyama, T., see Fujii, T., 181
Krämer, M., see Dabbert, D., 205
Kreis, M.E., Müller, M., Zittel, T.T., Glatzle, J., Grundy, D., Mediators of neuronal activation in the rat brain stem following intestinal anaphylaxis, 45
Kumlien, E., see Samuelsson, C., 185

Légrádi, G., see Fekete, C., 152
Laplanche, L., Kamenka, J.-M., Barbanel, G., The novel non-competitive *N*-methyl-D-aspartate antagonist gacyclidine blocks the glutamate-induced release of hydroxyl radicals in the striatum under conditions in which dizocilpine does not. A microdialysis study in rats, 49
Latorraca, S., see Gestri, D., 157
Lechan, R.M., see Fekete, C., 152
Lee, H.-P., see Choi, Y.-G., 173
Lee, J.-C., see Won, M.H., 119
Lestienne, F.G., see Ghafouri, M., 193
Libro, G., see de Tommaso, M., 57
Liguri, G., see Gestri, D., 157
Lindholm, D., see Samuelsson, C., 185

Mäder, M., see Dabbert, D., 205
Müller, D.J., see Block, W., 147

Müller, M., see Kreis, M.E., 45
Mackert, B.-M., Burghoff, M., Hiss, L.-H., Nordahn, M., Trahms, L., Curio, G., Non-invasive magnetoneurography for 3D-monitoring of human compound action current propagation in deep brachial plexus, 33
Magnaghi, V., see Petroni, A., 139
Maier, W., see Block, W., 147
Maier, W., see Majores, M., 224
Majores, M., Bagli, M., Papassotiropoulos, A., Schwab, S.G., Jessen, F., Rao, M.L., Maier, W., Heun, R., Allelic association between the D10S1423 marker and Alzheimer's disease in a German population, 224
Mamede-Rosa, M.L.N., see de Oliveira, R.M.W., 123
Maréchal, P.-A., see Boireau, A., 95
Martí, M.J., see Muñoz, E., 66
Massaro, A.M., see Gestri, D., 157
Matsumoto, T., see Takagi, H., 169
Matsumoto, Y., see Moon, C., 41
Matsushima, O., see Wang, X., 115
Matto, V., Vaarmann, A., Rudissaar, R., Pruus, K., Skrebuhhova-Malmros, T., Allikmets, L., Apomorphine-induced aggressive behaviour and post-mortem monoamine content in male Wistar rats, 131
Melcangi, R.C., see Petroni, A., 139
Meunier, M., see Boireau, A., 95
Mezey, E., see Charles, V., 29
Michel, O., see Hess, A., 72
Mihály, E., see Fekete, C., 152
Miwa, T., see Takahashi, S., 21
Mogi, M., Togari, A., Tanaka, K.-i., Ogawa, N., Ichinose, H., Nagatsu, T., Increase in level of tumor necrosis factor-α in 6-hydroxydopamine-lesioned striatum in rats is suppressed by immunosuppressant FK506 (tacrolimus), 165
Moon, C., Kim, S., Wie, M., Kim, H., Cheong, J., Park, J., Jee, Y., Tanuma, N., Matsumoto, Y., Shin, T., Increased expression of p53 and Bax in the spinal cords of rats with experimental autoimmune encephalomyelitis, 41
Mori, M., see Takagi, H., 169
Morishita, F., see Wang, X., 115
Moussaoui, S., see Boireau, A., 95
Mullan, M.J., see Crawford, F.C., 61
Muñoz, E., Pastor, P., Martí, M.J., Oliva, R., Tolosa, E., A new mutation in the parkin gene in a patient with atypical autosomal recessive juvenile parkinsonism, 66

Nagai, M., Wada, M., Usui, N., Tanaka, A., Hasebe, Y., Pleasant odors attenuate the blood pressure increase during rhythmic handgrip in humans, 227
Nagatsu, T., see Mogi, M., 165
Nakajima, Y., see Fujii, T., 181
Nordahn, M., see Mackert, B.-M., 33
Nowak, H., see Zwiener, U., 103
Nozawa, A., see Kakuda, T., 189

O'Callaghan, J.P., see Pennypacker, K., 1
Ogawa, N., see Mogi, M., 165
Oka, T., see Osako, Y., 25
Oliva, R., see Muñoz, E., 66
Orlacchio, A., see Gestri, D., 157
Osako, Y., Otsuka, T., Taniguchi, M., Oka, T., Kaba, H., Oxytocin depresses spontaneous γ aminobutyric acid-ergic inhibitory postsynaptic currents in cultured mitral cells of the rat olfactory bulb by a presynaptic mechanism, 25
Otokawa, M., see Hasegawa, H., 161
Otsuka, T., see Osako, Y., 25

Padovan, C.M., see de Oliveira, R.M.W., 123
Papassotiropoulos, A., see Majores, M., 224
Papini, N., see Petroni, A., 139
Park, J., see Moon, C., 41
Park, S.-K., see Won, M.H., 119
Pastor, P., see Muñoz, E., 66
Pennypacker, K., Eidizadeh, S., Kassed, C.A., O'Callaghan, J.P., Sanberg, P.R., Willing, A.E., Expression of fos-related antigen-2 in rat hippocampus after middle cerebral arterial occlusion, 1
Petroni, A., Papini, N., Blasevich, M., Magnaghi, V., Cavarretta, I., Galli, C., Melcangi, R.C., Testosterone metabolites in patients reduce the levels of very long chain fatty acids accumulated in X-adrenoleukodystrophic fibroblasts, 139
Philippu, A., see Singewald, N., 17
Polymeropoulos, M.H., see Charles, V., 29
Poon, P.W.F., Yu, P.P., Spectro-temporal receptive fields of midbrain auditory neurons in the rat obtained with frequency modulated stimulation, 9
Pruus, K., see Matto, V., 131
Puca, F., see de Tommaso, M., 57

Ramos, M.C., see Bullido, M.J.V., 213
Rand, W.M., see Fekete, C., 152
Rao, M.L., see Majores, M., 224
Ravindranath, V., see Annepu, J., 209
Reddy, P.H., see Charles, V., 29
Rempel, N., see Casper, D., 201
Ribeiro, J.A., see Cunha, R.A., 127
Richards, D., see Crawford, F.C., 61
Rietschel, M., see Block, W., 147
Rodríguez-Artalejo, F., see Bullido, M.J.V., 213
Ronne-Engström, E., see Samuelsson, C., 185
Rösner, S., see Dabbert, D., 205
Rudissaar, R., see Matto, V., 131

Sánchez-Criado, J.E., see González, D., 111
Saito, M., see Takagi, H., 169
Sakuma, K., see Wang, Y., 135
Samuelsson, C., Kumlien, E., Flink, R., Lindholm, D., Ronne-Engström, E., Decreased cortical levels of astrocytic glutamate transport protein GLT-1 in a rat model of posttraumatic epilepsy, 185
Sanberg, P.R., see Pennypacker, K., 1
Sastre, I., see Bullido, M.J.V., 213
Sato, R., see Takagi, H., 169
Satoh, K., Suzuki, N., Yokota, H., ADAMTS-4 (a disintegrin and metalloproteinase with thrombospondin motifs)-4 is transcriptionally induced in beta-amyloid treated rat astrocytes, 177
Sattelle, D.B., Harrison, J.B., Chen, H.H., Bai, D., Takeda, M., Immunocytochemical localization of putative γ-aminobutyric acid receptor subunits in the head ganglia of *Periplaneta americana* using an anti-RDL C-terminal antibody, 197
Schöll, U., see Dabbert, D., 205
Schild, H.H., see Block, W., 147
Schinka, J., see Crawford, F.C., 61
Schulze, T.G., see Block, W., 147
Schwab, S.G., see Majores, M., 224
Sciruicchio, V., see de Tommaso, M., 57
Sevush, S., see Crawford, F.C., 61

Shin, T., see Moon, C., 41
Shirokawa, T., Ishida, Y., Isobe, K.-i., Changes in electrophysiological properties of axon terminals of locus coeruleus neurons with age in F344 rat, 69
Singewald, N., Kouvelas, D., Kaehler, S.T., Sinner, C., Philippu, A., Peripheral chemoreceptor activation enhances 5-hydroxytryptamine release in the locus coeruleus of conscious rats, 17
Sinner, C., see Singewald, N., 17
Skrebuhhova-Malmros, T., see Matto, V., 131
Smith, C., see Sylvester, L., 13
Sorbi, S., see Gestri, D., 157
St. George-Hyslop, P.H., see Gestri, D., 157
Stennert, E., see Hess, A., 72
Suzuki, H., see Takagi, H., 169
Suzuki, N., see Satoh, K., 177
Sylvester, L., Kapron, C.M., Smith, C., In utero ethanol exposure decreases rapid eye movement sleep in female Sprague–Dawley rat offspring, 13

Tagle, D.A., see Charles, V., 29
Takagi, H., Sato, R., Mori, M., Matsumoto, T., Saito, M., Ito, E., Suzuki, H., Time-sharing contributions of A- and D-type K^+ channels to the integration of high-frequency sequential excitatory post synaptic potentials at a model dendrite in rats, 169
Takahashi, S., Miwa, T., Horikomi, K., Involvement of σ_1 receptors in methamphetamine-induced behavioral sensitization in rats, 21
Takeda, M., see Sattelle, D.B., 197
Takeuchi, S., see Watanabe, H., 53
Tanaka, A., see Nagai, M., 227
Tanaka, K.-i., see Mogi, M., 165
Tanaka, R., see Watanabe, H., 53
Taniguchi, M., see Osako, Y., 25
Tanuma, N., see Moon, C., 41
Tatro, J.B., see Fekete, C., 152
Tedde, A., see Gestri, D., 157
Tepest, R., see Block, W., 147
Togari, A., see Mogi, M., 165
Tolosa, E., see Muñoz, E., 66
Träber, F., see Block, W., 147
Trahms, L., see Mackert, B.-M., 33
Tumani, H., see Dabbert, D., 205

Unno, T., see Kakuda, T., 189
Usui, N., see Nagai, M., 227
Utsunomiya, K., see Kakuda, T., 189
Uysal, N., see Gönenç, S., 107

Vaarmann, A., see Matto, V., 131
Valdivieso, F., see Bullido, M.J., 213

Wada, M., see Nagai, M., 227
Wang, X., Morishita, F., Matsushima, O., Fujimoto, M., *Carassius* RFamide, a novel FMRFa-related peptide, is produced within the retina and involved in retinal information processing in cyprinid fish, 115
Wang, Y., Sakuma, K., Kakigi, R., The dynamic processes for word and picture encoding in the human brain as revealed by magnetoencephalography, 135
Watanabe, H., Abe, H., Takeuchi, S., Tanaka, R., Protective effect of microglial conditioning medium on neuronal damage induced by glutamate, 53
Weber, F., see Dabbert, D., 205
Weiss, K.R., see Klein, A.N., 37
Werner, P., see Casper, D., 201
Wie, M., see Moon, C., 41
Wilczak, N., see Zeinstra, E., 75
Willing, A.E., see Pennypacker, K., 1
Wirtshafter, D., A comparison of the patterns of striatal Fos-like immunoreactivity induced by various dopamine agonists in rats, 99
Wittekindt, C., see Hess, A., 72
Won, M.H., Kang, T.-C., Lee, J.-C., Choi, K.-Y., Park, S.-K., Jeong, Y.G., Jo, S.M., Age-related change of neuropeptide Y-immunoreactive neurons in the rat main olfactory bulb, 119

Yanase, H., see Kakuda, T., 189
Yaparpalvi, U., see Casper, D., 201
Yasumatsu, M., see Hasegawa, H., 161
Yazawa, T., see Hasegawa, H., 161
Yokota, H., see Satoh, K., 177
Young, T.A., see Charles, V., 29
Yu, P.P., see Poon, P.W.F., 9

Zeinstra, E., Wilczak, N., De Keyser, J., [^{3}H]dihydroalprenolol binding to beta adrenergic receptors in multiple sclerosis brain, 75
Zittel, T.T., see Kreis, M.E., 45
Zwiener, U., Eiselt, M., Gießler, F., Nowak, H., Relations between early prespike magnetic field changes, interictal discharges, and return to basal activity in the neocortex of rabbits, 103